

An echo of an exciting light pulse in quantum wells

I. G. Lang, L. I. Korovin

A. F. Ioffe Physical-Technical Institute, Russian Academy of Sciences, 194021 St. Petersburg, Russia

D. A. Contreras-Solorio, S. T. Pavlov[†]

Escuela de Física de la UAZ, Apartado Postal C-580, 98060 Zacatecas, Zac., Mexico

(February 1, 2008)

The non-sinusoidal character oscillations appear in the transmitted, reflected and absorbed light fluxes when light pulses irradiate a semiconductor quantum well (QW), containing a large number of the equidistant energy levels of electronic excitations. A damping echo of the exciting pulse appears through the time intervals $2\pi\hbar/\Delta E$ in the case of the very short light pulses $\gamma_l^{-1} \ll \hbar/\Delta E$.

A row of theoretical and experimental investigations is devoted to the elaborate study of the electronic excitations in the bulk crystals and semiconductor QWs with the help of the time resolved scattering (TRS), because just the existence of the discrete energy levels determines the most interesting results obtainable by the TRS. It is well known that a couple of the closely disposed energy levels demonstrate a new effect: The sinusoidal beatings appear in reflected and transited pulses on a frequency corresponding to the energy distance between the energy levels [1].

We investigate theoretically below reflection and absorption of light pulses, irradiating semiconductor QWs, having a large number of equidistant energy levels. Let us suppose that a QW with a system of equidistant discrete energy levels is irradiated by a light pulse with a carrier frequency ω_l . Let us suppose that the frequency ω_l is in the resonance with one of energy levels.

Then two variants are possible: first, one may neglect influence of the rest energy levels; second, one have to take into account a few neighbor energy levels. The first variant is realized, for instance, for a symmetrical pulse under condition $\gamma_l \ll \Delta\omega$, where $\Delta\omega$ is the distance between the neighbor levels, γ_l is the parameter characterizing the pulse duration. It is shown [2] that under condition $\gamma_r \geq \gamma$, where $\gamma_r(\gamma)$ is the radiative (non-radiative) broadening of an electronic excitation, the transmitting pulse profile changes drastically. The lifetimes of the electron-hole pairs (EHP) in a QW in a quantizing magnetic field have been calculated in [3,4]. The second variant with many equidistant energy levels has been considered in [3,5], where a ladder-like structure of a reflected and transmitted pulses has been predicted.

We consider here both symmetrical and asymmetrical pulses, demonstrate the results for the different γ_l and show that very short pulses create "an echo" in reflected and transmitted light.

Let us consider a QW with a system of equidistant energy levels

$$\omega_\rho = \rho\Omega_\mu, \quad \rho = 0, 1, 2, \dots \quad (1)$$

The real significance of the designation Ω_μ will be given below. We believe that the QW's width is much smaller than the light wavelength $d \ll c/(n\omega_l)$, where n is the refraction index outside the QW. The excitation ρ is characterized by the radiative $\gamma_{r\rho}$ and non-radiative broadenings.

We suppose that the light pulse incidents perpendicular the QW's plane xy from the left (from the negative z) and its electric field is

$$\mathbf{E}_0(\mathbf{z}, t) = \mathbf{E}_0 \mathbf{e}_l \exp(-i\omega_l t) \{ \Theta(p) \exp(-\gamma_{l1} p/2) + [1 - \Theta(p)] \exp(\gamma_{l2} p/2) \} + c.c., \quad (2)$$

where E_0 is the real amplitude, $\mathbf{e}_l$ is the circular polarization vector, $p = zn/c$, $\Theta(p)$ is the Heaviside function. We have for a symmetrical pulse

$$\gamma_{l1} = \gamma_{l2} = \gamma_l, \quad (3)$$

and for an pulse with a sharp front

$$\gamma_{l1} = \gamma_l, \quad \gamma_{l2} \rightarrow \infty. \quad (4)$$

The electric fields on the left (right) of the QW and corresponding time-dependent energy fluxes are calculated according to the methods of [2–6]. We demonstrate some results obtained for the following parameters

$$\gamma_{r\rho} = \gamma_r, \quad \gamma_\rho = \gamma, \quad \gamma_r \ll \gamma \ll \gamma_l, \quad \gamma \ll \Omega_\mu, \quad (5)$$

and for an arbitrary interrelation of γ_l and Ω_μ . Let us consider the energy fluxes under conditions

$$p \gg \gamma_l^{-1}, \quad s \gg \gamma_l^{-1}, \quad p \ll \gamma_r^{-1}, \quad s \ll \gamma_r^{-1}, \quad (6)$$

where $s = t + zn/c$ is the variable for the reflected energy flux. Then the contributions into the electric fields, containing the factors $\exp(-\gamma_l p/2)$ or $\exp(-\gamma_l s/2)$, become negligible. We obtain, that only those contributions in induced fields are essential which are proportional to $\exp(-\gamma_r p/2)$ or $\exp(-\gamma_r s/2)$. The existence of

such contributions is proved in many investigations. A picture of induced fields outside a QW is symmetrical, since only narrow QWs are considered. It follows from this fact that the transmitted and reflected fluxes are equal in absolute values and the absorbed flux equals to the doubled transmitted (or reflected) flux with the opposite sign. A negative absorption means that the QW gives back a stored energy, irradiating it symmetrically by two fluxes on the left and on the right.

The reflected energy flux results in

$$\mathcal{R}(s) = 4(\gamma_r/\gamma_l)^2 e^{-\gamma s} Y_{\Omega_l, G_l}(S), \quad (7)$$

where $\mathcal{R}(s)$ is the module of the reflected flux in units $cE_0^2/(2\pi n)$, Y_{Ω_l, G_l} is the dimensionless periodical function of the variable $S = \Omega_\mu s$ with the period 2π , depending from the parameters

$$\Omega_l = \omega_l/\Omega_\mu, \quad G_l = \gamma_l/\Omega_l. \quad (8)$$

The factor $\exp(-\gamma s)$ determines damping of the reflected flux. The function Y is periodical on s with the period $2\pi/\Omega_\mu$. These oscillations are not sinusoidal.

We calculated the functions $Y_{\Omega_l, G_l}(S)$ and showed, that the results are essentially dependent on the parameter G_l . For $G_l \ll 1$ the periodical vibrations of the intensities of the reflected and transmitted energy fluxes are very small in amplitudes. The values $\Omega_l = 0, 1, 2, \dots$ correspond to the resonance of the frequency ω_l with one of the energy levels. A small detuning of the frequency ω_l results into the drastic dropping of the energy fluxes.

The function $Y_{\Omega_l, G_l}^{sym}(S)$ is represented in Figs. 1, 2 inside of one period.

The upper index means a symmetrical exciting pulse. The function $Y_{\Omega_l, G_l}^{sym}(S)$ is symmetrical relatively the substitution S by $2\pi - S$. Fig. 1 corresponds to the value G_l and to the set $\Omega_l = 0; 0.1; 0.5; 1; 1.5; 2$. The vibration amplitude in Fig.1 reaches the value 1.5-2. For $\Omega_l = 0.5$ and $\Omega_l = 1.5$, i. e. in the case when the frequency is disposed between the levels $n = 0$ and $n = 1$ and between the levels $n = 1$ and $n = 2$, respectively, the curves touch the abscissas axis in the point $S = \pi$.

Fig. 2 demonstrates the echo of the exciting symmetrical pulse. It corresponds to the large value $G_l = 5$ and to the set $\Omega_l = 0; 0.1; 0.5; 1$. The function $Y^{sym}(S)$

in the points $s = 0$ $s = 2\pi$ increases sharply in comparison to corresponding values in the Fig. 1, reaching 30, but they are very small in the interval $s \gg G_l^{-1}$, $(2\pi - S) \gg G_l^{-1}$. Thus, the periodical function $Y^{sym}(S)$ represents a sequence of short pulses, the duration of each is of the order G_l^{-1} and disposed with the intervals 2π . Applying Eq. (7), one finds, that at $\gamma_l \gg \Omega_\mu$ some echo of the exciting pulse has to be seen in the reflected energy flux, damping as $\exp(-\gamma s)$ with the interval $2\pi/\Omega_\mu$. Such echo must be also in the transmitted energy flux. Our results show that in the region of

values of Ω_l from 0 up to several units, the pulse replica form repeats. In a resonance ω_l with one of upper levels with index $\rho_0 \gg 1$ the form of the repeating pulses (echo) coincides with the exciting pulse form. The repeating pulses contain the small factor

$$\delta = \pi^2(\gamma_r/\Omega_\mu)^2 \exp(-\gamma s). \quad (9)$$

The theory is applicable for the narrow QWs in a strong magnetic field perpendicular to the QW's plane xy . Then the equidistant levels correspond to the EHP with different Landau levels and fixed size quantized energy levels of electrons (holes). $\Omega_\mu = |e|H/\mu c$ is the cyclotron frequency, corresponding to the reduced mass $\mu = m_e m_h / (m_e + m_h)$, where $m_e(m_h)$ is the electron (hole) effective mass. The energy level equidistance is small if

$$d \ll a, \quad a_H \ll a, \quad (10)$$

where a is the radius of the Wannier=Mott exciton in a zero magnetic field, $a_H = (\hbar c / |e|H)^{1/2}$ is the magnetic length, i. e. for the narrow QWs and strong magnetic fields. We suppose that the band's non-parabolicity is small inside the vicinity of the band's extremums.

Acknowledgements

S.T.P thanks the Zacatecas University and the National Council of Science and Technology (CONACyT) of Mexico for the financial support and hospitality. D.A.C.S. thanks CONACyT (27736-E) for the financial support. This work has been partially supported by the Russian Foundation for Basic Research and by the Program "Solid State Nanostructures Physics".

[†] On leave from the P. N. Lebedev Physical Institute, 117924 Moscow, Russia.

- [1] H. Stolz, Time - Resolved Light Scattering from Excitons, Springer Tracts in Modern Physics, Springer, Berlin, 1994.
- [2] L. I. Korovin, I. G. Lang, A. Contreras Solorio and S. T. Pavlov, Fiz. Tverd. Tela, **42**, issue 12, p. 119 (2000); LANL archiv/cond-mat/0006364.
- [3] I. G. Lang, V. I. Belitsky, M. Cardona, Phys. stat. sol. (a), **164** 307 (1997).
- [4] I. G. Lang, L. I. Korovin, A. Contreras Solorio and S. T. Pavlov, LANL archiv/cond-mat/0001248 .
- [5] I. G. Lang, V. I. Belitsky. Physics Letters A, 245 , 329 (1998).
- [6] D. A. Contreras Solorio, S. T. Pavlov, , L. I. Korovin and I. G. Lang, Phys. Rev. B, **62**, N 24, pp. 16815-16819 (2000), LANL archiv/cond-mat/0002229 .

- [7] I. V. Lerner, Yu. E. Lozovik, Zh. Eksp. i Teor. Phys., **78**, 1167 (1980).

FIG. 1. The function $Y_{\Omega_l, G_l}^{sim}(S)$, corresponding to the periodical factor in the value of the reflected energy flux when a QW is irradiated by the symmetrical light pulse. The pulse duration $\gamma_l^{-1} = \Omega_\mu^{-1}$.

FIG. 2. Same as Fig.1 for the very small value γ_l^{-1} , when the echo of the exciting pulse appears.

This figure "fig1.JPG" is available in "JPG" format from:

<http://arXiv.org/ps/cond-mat/0105198v1>

This figure "fig2.JPG" is available in "JPG" format from:

<http://arXiv.org/ps/cond-mat/0105198v1>